



# Rodilon®


TRIO

## SINGLE-FEED RODENTICIDE

For Professional Use Only

A grain bait containing Difethialone (0.0025% w/w) for use as a rodenticide by professionals for the control of rats and mice indoors for the protection of public health, stored products and materials.

Kills Mice and Rats.



## TRIO

- Not for amateur sale
- Ideal for mice
- Active ingredient locked in

Rodilon® contains 0.0025% w/w difethialone.

Harmful to aquatic life with long lasting harmful effects.

Keep out of reach of children.

Read label before use.

Keep/Store away from food drink and animal feedingstuffs.

Do not eat, drink or smoke when using this product.

Avoid release to the environment.

Wear protective gloves.

IF SWALLOWED: Immediately call a poison centre or doctor/physician.

Store locked up in a closed container.

Dispose of contents/container in accordance with national regulations.

IE/BPA 70182

## 800 g e



# Rodilon®

TRIO

# Rodion® Trio

A grain bait containing Difethialone (0.0025% w/w) for use as a rodenticide by professionals for the control of rats and mice indoors for the protection of public health, stored products and materials. Rodion Trio is a highly potent anticoagulant mixed grain bait for indoor use. It contains the newest active ingredient (Difethialone 0.0025% w/w) introduced to rodent control. The mixed grains (Oat, sunflower and maize) used in this formulation have been chosen from the most palatable grains known to increase rodent consumption, even in the most difficult treatment environments. This, combined with the "turbo impregnation" technology used in production, delivers Rodion Trio its extreme palatability and overall superior performance which makes it the ideal product for mice control. Turbo impregnation: Each grain is coated and impregnated through to its core. This gives the following benefits over other rodenticides:

- Enhanced mice control (Mice remove the exterior of the grain and eat the middle).
- Significantly increased rat control due to more consistent coverage of each grain.
- Minimal dust, reducing exposure to the user.

Rodion Trio kills House Mice (*Mus musculus*), Norway/Brown Rats (*Rattus norvegicus*) and Black Rats (*Rattus rattus*), including all known resistant strains to other anti-coagulants baits.

## DIRECTIONS FOR USE:

Rodion Trio is a highly palatable mixed grain bait based on the latest active ingredient Difethialone and can be used indoors only, in areas including agricultural, domestic, commercial, food factories, public services, municipal and industrial buildings. The effects on rodents can be seen 3 to 5 days after the first consumption of the bait. Rats may take several days before starting to feed because of their neophobic behaviour (fear of anything new). Rats and mice can consume a lethal dose in one feed. Place Rodion Trio where it is inaccessible to children, animals and birds. Bait must be securely deposited in a baiting station or other coverings so as to minimize the risk of consumption by other animals or children. Where possible secure baits so they cannot be dragged away. The red colouring of the grains shows in the rodent excrement indicating bait has been taken. Wash hands after use of the product, before eating, drinking and smoking.

## Application:

Indoors in urban and agricultural buildings.

## Baiting for Mice:

Place up to 50 g of bait in tamper resistant bait stations or covered bait points spaced 2-3 m apart (1-1.5 m apart in high infestation areas) in areas where mice are active. Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation, (e.g. wherever damage, droppings or evidence of mice activity has been found). Mice feed sporadically and will consume small quantities from numerous feeding points. Eliminate if possible all alternative foods.

## Baiting for Rats:

Place up to 200 g of bait in tamper resistant bait stations or covered bait points spaced 8-10 m apart (4-5 m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation, (e.g. wherever damage, tracks, droppings, nests and along rodent runs). Rats will consume from a single feeding point. Eliminate if possible all alternative foods.

## For Rats and Mice:

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped.

Authorisation holder :  
Lipilatch S.A.S - Bommel  
Rue de l'Étoile du Casse  
(France)  
Phone: +33 553 693570

For 24 hour emergency information contact  
Bayer CropScience Ltd.  
Freephone: 1800 409399  
or nearest National Poisons Information Centre  
Rodion® is a registered trademark of Bayer.  
©Copyright of Bayer 2015 - All rights reserved  
Bitrex® is a registered trademark  
of Macfarlan Smith, Edinburgh

Marketing Company:  
Bayer CropScience Limited,  
230 Cambridge Science Park,  
Milton Road, Cambridge, CB4 0WB.  
Tel No. 08000 1214 9451  
www.pestcontrol-expert.com  
for MSDS and larger label

## Follow the baiting strategy as recommended below :

Baiting for:	Level of infestation	Recommended Application rates	Inspection and replacing
Mice	High infestation	up to 50 g per baiting point every 1 to 1.5 meters	Inspect frequently especially in first 5-10 days, and as regularly as consumption persists (if possible every week). At each inspection, replace any bait that has been consumed. Replace any contaminated or spoilt bait. Maintain a constant supply of fresh bait until signs of rodent activity cease.
	Low infestation	up to 50 g per baiting point every 2 to 3 meters	
Rats	High infestation	up to 200 g per baiting point every 4 to 5 meters	
	Low infestation	up to 200 g per baiting point every 8 to 10 meters	

## STATUTORY CONDITIONS OF APPROVAL RELATING TO USE.

For professional use only. Use Biocides Safely. Always Read the label and product information before use.

For use as a rodenticide only. To avoid risks to human health and the environment comply with instructions for use. Use biocides safely and sustainably, it is illegal to use this product for uses in a manner other than that presented on the label. Harmful to aquatic life and may cause lasting harmful effects to aquatic life. Harmful if swallowed. Harmful to wildlife. Keep locked up and out of reach of children. Keep away from food, drink and animal feed stuffs. When using, do not eat, drink or smoke. This material and its container must be disposed of in a safe way. Do not empty into drains. Wear suitable gloves. If swallowed, seek medical advice immediately and show this container or label. Avoid release into the environment. Refer to the safety data sheet. Use bait containers clearly marked "poison" at all surface baiting points. Remove all remains of bait and dead rodents during and after treatment and dispose of safely. Apply only in positions inaccessible to children, domesticated animals and pets (particularly cats and dogs and pigs). When the product is used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in the case of poisoning, must be made available alongside the baits.

IE/BA 70182

## FOLLOW-UP:

Dead rodent bodies, remains of unused baits or any fragments of bait found away from the bait station must be collected during all operations to minimize the risk of consumption and poisoning to children, companion animals and other non-target animals.

## Precautions:

Keep out of reach of children. Read label before use. Use Biocides safely and sustainably. It is illegal to use this product for uses or in a manner other than prescribed on this label. NOT for amateur sale.

## In case of accidental poisoning:

If swallowed immediately call The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166) or a doctor/physician, and refer to the label. Because of anti vitamin K properties of the active ingredient, absorption can inhibit blood clotting and cause hemorrhagic syndrome- bruising easily, nose or gum bleeds, blood in stools or urine, excessive bleeding from minor cuts or abrasions. Note that poisoning symptoms may develop over the course of several days. Note to Physician: Antidote - Vitamin K1 under medical supervision.

## FIRST AID:

In case of accident, suspected exposure or if you feel unwell seek medical advice immediately (show the label where possible). In case of contact with skin, wash soap and water. Remove and launder contaminated clothing. In case of contact with eyes remove contact lenses if present and rinse the eye slowly and gently with water for 15-20 minutes, seek medical help immediately.

## Resistance Information:

The resistance status of the rodent population should be taken into account when considering the choice of rodenticide to be used. This requirement is to help minimise resistance to difethialone. Where resistance to difethialone has been shown or is suspected resistance management strategies should be employed.

## Storage and Disposal:

Do not contaminate water, food or feed by storage or disposal. This is a non refillable container. Do not decant into unlabelled containers. Store locked up in the original container in a cool, dry, well ventilated place. Do not reuse or refill. This material and its container must be disposed of safely. Dispose of packaging, remains of unused products and dead rodents in accordance with local regulations, if required consult a professional waste operator or local authority.

IE80862903C - ARTICLE 80572999 - RA2

Date of expiry: see the pack  
Batch number: see the pack

