

VERTOX

CONTACT GEL



300g

A READY-FOR-USE CONTACT GEL RODENTICIDE FOR THE CONTROL OF MICE INDOORS.
CONTAINS BRODIFACOUM 0.005% W/W AND DENATONIUM BENZOATE 0.001% W/W

DIRECTIONS FOR USE

Place contact gel where there are signs of mouse activity, like fresh droppings, inside or near nesting area, runways and mouse feeding places. Place contact gel points throughout the infested area. Pay special attention to non-target risks both from direct access to the gel and from the possibility of secondary poisoning from dead mice. After use, seal mastic gun securely and keep in a safe place out of reach of children and animals.

For mouse infestations use contact gel points of up to 15 g. Place gel points 5 m apart reducing to 2 m in cases of high infestation.

Apply the contact gel as a 'zig-zag' strip or thin layer on pieces of plastic or card and fix in areas likely to be walked through by mice. Contact gel that adheres to the feet and fur of the mice will be removed during normal grooming behaviour and ingested. Protect contact gel points from non-target animals and from water, preferably using commercially available bait stations. Where these are not available, place the contact gel on bait trays or on plastic strips inside pieces of drainage pipe or under sections of slate, board or corrugated iron. Contact gel can be placed on strips of insulation tape or paper tape fixed to overhead pipe-ways and ductwork. Tape can be easily removed after treatment is concluded. Ensure that contact gel cannot fall or become displaced onto sensitive areas during use. Contact gel can be applied to holes in walls and partitions through which mice run. Make frequent inspections of the contact gel points during the first 10-14 days and replace any contact gel removed by rodents or that has been damaged by water or contaminated by dirt. If all the contact gel has been removed from certain areas, increase the quantity of contact gel by placing more gel points. Do not increase the gel point size. Precautions should be taken to prevent domestic animals from having access to the contact gel. The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used.

This product must be supplied with an 'advice to householder leaflet' containing details of the product and active substance used, that the product is an anticoagulant and a precaution to prevent access to bait by children and domesticated animals particularly dogs and birds.



PelGar International Ltd.

Unit 13 Newman Lane, Alton,
Hampshire, GU34 2QR.
Tel - 01420 80744
Email - info@pelgar.co.uk www.pelgar.co.uk

STATUTORY CONDITIONS RELATING TO USE

FOR USE ONLY IN MOUSE CONTROL BY PROFESSIONAL OPERATIONS. FOR INDOOR USE ONLY. Apply as necessary. Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.

READ ALL PRECAUTIONS BEFORE USE. HSE No. 9296

PRECAUTIONS

USE ONLY AS DESCRIBED ON THE LABEL.

*The (COSHH) Control of Substances Hazardous to Health Regulations 2002 may apply to the use of this product at work. *Guidance on the safe use of rodenticides is provided in leaflets AIS 31 and MISC 515 at <http://www.hse.gov.uk/pubns/biopestindex.htm>.

*Do not place contact gel where food, feed or water could become contaminated. *PREVENT ACCESS TO CONTACT GEL by children, birds and non-target animals particularly dogs, cats, pigs and poultry. *Avoid excessive contamination of coveralls. *Search for and remove rodent bodies at frequent intervals, ideally each time you inspect the bait but at least every 2-3 days during the first 14 days of contact gel placement. Collect and dispose of the remains of contact gel and any remaining rodent bodies after treatment. You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance contact the Environment Agency or your local Scottish Environmental Protection Agency (SEPA) office.

*HARMFUL TO WILDLIFE. *Avoid contact with skin. *Wild mammals and birds may be at greater risk if this product is not used in accordance with the label. *Keep away from food, drink and animal feedingstuffs. *Brodifacoum is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin time not less than eighteen hours after consumption. If elevated, administer Vitamin K1 until prothrombin time normalises. Continue determination of prothrombin time for two weeks after withdrawal of the antidote and resume treatment if elevation occurs in that time. *IF YOU FEEL UNWELL, seek medical advice (show the label where possible). *WASH HANDS AND EXPOSED SKIN before meals and after use. *Keep only in original container. *Keep in a safe place. *Empty container completely and dispose of safely.

To avoid risks to man and the environment, comply with the instructions for used.

Protect from extremes of temperature. Store in original sales pack between +5°C and +25°C.

VERTOX

CONTACT GEL



300g

A READY-FOR-USE CONTACT GEL RODENTICIDE FOR THE CONTROL OF MICE INDOORS.
CONTAINS BRODIFACOUM 0.005% W/W AND DENATONIUM BENZOATE 0.001% W/W

DIRECTIONS FOR USE

Place contact gel where there are signs of mouse activity, like fresh droppings, inside or near nesting area, runways and mouse feeding places. Place contact gel points throughout the infested area. Pay special attention to non-target risks both from direct access to the gel and from the possibility of secondary poisoning from dead mice. After use, seal mastic gun securely and keep in a safe place out of reach of children and animals.

For mouse infestations use contact gel points of up to 15 g. Place gel points 5 m apart reducing to 2 m in cases of high infestation.

Apply the contact gel as a 'zig-zag' strip or thin layer on pieces of plastic or card and fix in areas likely to be walked through by mice. Contact gel that adheres to the feet and fur of the mice will be removed during normal grooming behaviour and ingested. Protect contact gel points from non-target animals and from water, preferably using commercially available bait stations. Where these are not available, place the contact gel on bait trays or on plastic strips inside pieces of drainage pipe or under sections of slate, board or corrugated iron. Contact gel can be placed on strips of insulation tape or paper tape fixed to overhead pipe-ways and ductwork. Tape can be easily removed after treatment is concluded. Ensure that contact gel cannot fall or become displaced onto sensitive areas during use. Contact gel can be applied to holes in walls and partitions through which mice run. Make frequent inspections of the contact gel points during the first 10-14 days and replace any contact gel removed by rodents or that has been damaged by water or contaminated by dirt. If all the contact gel has been removed from certain areas, increase the quantity of contact gel by placing more gel points. Do not increase the gel point size. Precautions should be taken to prevent domestic animals from having access to the contact gel. The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used.

This product must be supplied with an 'advice to householder leaflet' containing details of the product and active substance used, that the product is an anticoagulant and a precaution to prevent access to bait by children and domesticated animals particularly dogs and birds.



PelGar International Ltd.

Unit 13 Newman Lane, Alton,
Hampshire, GU34 2QR.
Tel - 01420 80744
Email - info@pelgar.co.uk www.pelgar.co.uk

STATUTORY CONDITIONS RELATING TO USE

FOR USE ONLY IN MOUSE CONTROL BY PROFESSIONAL OPERATIONS. FOR INDOOR USE ONLY. Apply as necessary. Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.

READ ALL PRECAUTIONS BEFORE USE. HSE No. 9296

PRECAUTIONS

USE ONLY AS DESCRIBED ON THE LABEL.

*The (COSHH) Control of Substances Hazardous to Health Regulations 2002 may apply to the use of this product at work. *Guidance on the safe use of rodenticides is provided in leaflets AIS 31 and MISC 515 at <http://www.hse.gov.uk/pubns/biopestindex.htm>.

*Do not place contact gel where food, feed or water could become contaminated. *PREVENT ACCESS TO CONTACT GEL by children, birds and non-target animals particularly dogs, cats, pigs and poultry. *Avoid excessive contamination of coveralls. *Search for and remove rodent bodies at frequent intervals, ideally each time you inspect the bait but at least every 2-3 days during the first 14 days of contact gel placement. Collect and dispose of the remains of contact gel and any remaining rodent bodies after treatment. You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance contact the Environment Agency or your local Scottish Environmental Protection Agency (SEPA) office.

*HARMFUL TO WILDLIFE. *Avoid contact with skin. *Wild mammals and birds may be at greater risk if this product is not used in accordance with the label. *Keep away from food, drink and animal feedingstuffs. *Brodifacoum is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin time not less than eighteen hours after consumption. If elevated, administer Vitamin K1 until prothrombin time normalises. Continue determination of prothrombin time for two weeks after withdrawal of the antidote and resume treatment if elevation occurs in that time. *IF YOU FEEL UNWELL, seek medical advice (show the label where possible). *WASH HANDS AND EXPOSED SKIN before meals and after use. *Keep only in original container. *Keep in a safe place. *Empty container completely and dispose of safely.

To avoid risks to man and the environment, comply with the instructions for used.

Protect from extremes of temperature. Store in original sales pack between +5°C and +25°C.